

National Aeronautics and Space Administration  
Goddard Space Flight Center  
Contract No. NAS-5-3760

ST - CC - 10306

NASA TT F-9655

REGULITE FORM 902	N65-19732	
	(ACCESSION NUMBER)	(THRU)
	5	(CODE)
	(PAGES)	13
	(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

MEASUREMENT OF THE RADIOACTIVITY INDUCED BY COSMIC  
RADIATION IN THE BOGOU METEORITE

by

Daniel Nordemann

[ FRANCE ]

GPO PRICE \$ \_\_\_\_\_

OTS PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) \$1.00

Microfiche (MF) \$0.50

17 MARCH 1965

MEASUREMENT OF THE RADIOACTIVITY INDUCED BY COSMIC  
RADIATION IN THE BOGOU METEORITE

Comptes-Rendus de l'Académie  
des Sciences.- Météorites (Gr.10)  
T. 259, pp. 3581-3, Paris, 1964

by Daniel Nordemann  
& Jacques Tobailem

SUMMARY

The activities of  $\gamma$  - emitters contained in a sample of the Bogou meteorite, weighing 0.185 kg have been measured by quantitative  $\gamma$  - spectrometry of weak proper motion.

\* \* \*

An iron meteorite with mass of 8.8 kg fell near Dougou (Volta) on 14 August 1962 at 10 00 hours.

Our laboratory received a sample weighing 0.185 kg, in a form of thin plate (6 mm) cut in the mass of meteorite (see Fig.1). A first measurement was made 303 days after the fall and led to the evidence of an activity due to manganese 54 of  $750 \pm 75 \text{ dpm} \cdot \text{kg}^{-1}$  [1].

A subsequent measurement, effected by Mabuchi over the same sample, but with a different standardization method, gave the value of  $550 \pm 55 \text{ dpm} \cdot \text{kg}^{-1}$  [2].

We took over anew the study of that sample, using this time the  $\gamma$  - spectrometry, with new facilities permitting to obtain a background noise, weaker and more steady.

We have effected a series of measurements of long duration, up to 160 hours, spread between 680 and 760 days after the day of fall. Naturally, this time interval constitutes a disadvantage because of short-lived radionuclides (period  $\leq 1$  year), but, otherwise we may

---

\* Mesure de la radioactivité induite par le rayonnement cosmique dans la météorite de Bogou.

hope to detect more easily long-life radionuclides (sodium 22, cobalt 60, titanium 44, aluminum 26 etc.), which are much less concealed by the short-life radionuclides that have decreased.

The installation's calibration for these measurements was made by means of dummies simulating the meteorite sample.

Three dummies comprised calibrated sources of manganese 54, cobalt 60 and potassium 40. These models were materialized by interposing two plane sources, of homogenous distribution, between three stainless steel plates.

The following results, reduced to the date of the fall, have been obtained:

- <sup>54</sup> Mn	(period of 313.5 days)	[3] : 483 ± 40 dpm.kg <sup>-1</sup> ;
- <sup>22</sup> Na	( " 2.58 years)	[4] : 0 ± 4 " ;
- <sup>60</sup> Co	( " 5.27 years)	[5] : 14 ± 7 " ;
- <sup>44</sup> Ti	( " 200.0 years)	[6] : 3.8 ± 3 " ;
- <sup>26</sup> Al	( " 738,000 years)	[7] : 4.3 ± 3 " .

The value obtained for the activity of manganese 54 in the Bogou meteorite is in the vicinity of the values already published, that is  $485^{+30}_{-85}$  dpm.kg<sup>-1</sup> [8];  $439 \pm 22$  dpm.kg<sup>-1</sup> [9]. It is to be compared with those indicated for Aroos iron meteorite (fall of 24 November 1959) :  $470 \pm 47$  dpm.kg<sup>-1</sup> [10];  $425 \pm 40$  dpm.kg<sup>-1</sup> [11], and also with that obtained by Honda [12] by having the iron bombarded by 3 GeV protons : 440 dpm.kg<sup>-1</sup>.

The value of activity in cobalt 60 is sensibly in agreement with those indicated by other authors :  $\leq 7 \pm 4$  dpm.kg<sup>-1</sup> [9],  $\leq 13$  dpm.kg<sup>-1</sup> [8]. As to the activity in titanium 44 in the Bogou meteorite, no value has been published to-date.

The activity measured for aluminum 26 is in agreement with the value obtained by Shedlovskiy [13] after chemical separation, which is  $3.81 \pm 0.09$  dpm.kg<sup>-1</sup>.

In the iron meteorites, the principal formation modes of these nuclides, under the effect of cosmic radiation, are :

../..

- $^{54}\text{Mn}$  : spallation of iron by rapid secondary neutrons,  
 $^{54}\text{Fe}(n,p)^{54}\text{Mn}$  and  $^{56}\text{Fe}(n,p2n)^{54}\text{Mn}$ ;
- $^{60}\text{Co}$  : capture of secondary thermal neutrons by cobalt,  
 $^{59}\text{Co}(n,\gamma)^{60}\text{Co}$ ;
- $^{44}\text{Ti}$  : spallation of iron,  $^{56}\text{Fe}(p,5p8n)^{44}\text{Ti}$ ;
- $^{26}\text{Al}$  : spallation of iron,  $^{56}\text{Fe}(p,14p17n)^{26}\text{Al}$ .

Thus, the activity of Mn 54 must be influenced by the size of the meteorite and by the position of the sample inside. This explains the results' grouping as regards manganese 54 for all the various samples of a same meteorite and for meteorites having been subject to a comparable irradiation in space.




Fig. 1. - The Bogou Meteorite.

The hatched portion indicates the position of the 0.185 kg sample

To the contrary, the activity in cobalt 60 is a function of the shape of meteorite sample and of its position because of neutron production and thermalization [14]. On a meteorite of the size of that of Bogou the production of cobalt 60 at the center is more important than at the edges. This can explain the fact, that our measurement, made with the detector near center, gives an activity twice as great as that obtained

by Rowe, Anderson and Van Dilla [9], who analyzed a much more voluminous sample (8.4 kg), where the edges played a prominent part.

\*\*\* THE END \*\*\*

#### REFERENCES

- [1].- Y. YOKOHAMA & J. LABEYRIE., Nature, 201, p.809, 1964.
- [2].- H. MABUCHI.- Personal communication.
- [3].- E. I. WYATT, S. A. REYNOLDS, T. H. HANDLEY, W. S. LYON & H. A. PARKER.-  
Nuclear Sc. & Engineering, 11, 74, 1961.
- [4].- W. F. MERRITT, P. J. CAMPION & R. C. HAWKINGS.- Can.J.Phys. 35, 16, 1957.
- [5].- J. TOBAILEM.- Comptes-Rendus, 233, p.1360, 1951.
- [6].- M. MONDA & D. LAL. Nuclear Physics, 51, p.363, 1964.
- [7].- R. A. RIGHTMIRE, T. P. KOHMAN & H. HINTERBERGER.- Z. Naturforsch.,  
13a, p.847, 1958.
- [8].- J. P. UNIK, D. J. HENDERSON & J. R. HUIZENGA.- Geochimica & Cosmochimica  
Acta, 28, p.593, 1964.
- [9].- M. W. ROWE, E. C. ANDERSON & M. A. VAN DILLA.- J.Geophys.Res. 69, 1964
- [10].- M. HONDA & J. R. ARNOLD.- Geochimica and Cosmochimica Acta 23, 1961.
- [11].- M. W. ROWE, M. A. VAN DILLA & E. C. ANDERSON.- Ib., 27, p.1003, 1963
- [12].- M. HONDA.- J. Geophys. Res, 67, p.4847, 1962.
- [13].- J. P. SHEDLOVSKY.- Carnegie Inst.of Techn., Progress Rep. 2 B 5, p.86  
1962 - 1963
- [14].- P. EBERHARDT, J. GEISS & H. LUTZ., Earth Science  
& Meteorites, No.Holl.P.Co. 1963

---

(Centre d'Études Nucléaires de Saclay,  
Service d'Électronique physique,  
et Centre des Faibles Radioactivités du C. N. R. S.,  
Gif-sur-Yvette, Seine-et-Oise.)

---

CONTRACT No. NAS-5-3760

Consultants & Designers, Inc.  
Arlington, Virginia

Translated by ANDRE L. BRICHANT  
on 17 March 1965.

DISTRIBUTIONGODDARD SPACE F.C.

600 TOWNSEND  
STROUD  
610 MEREDITH  
611 McDONALD  
ABRAHAM  
BOLDT  
612 HEPPNER  
NESS  
613 KUPPERIAN [3]  
614 LINDSAY  
WHITE  
615 BOURDEAU  
BAUER  
STONE  
640 HESS [3]  
O'KEEFE  
660 GI for SS [5]  
252 LIBRARY [3]  
256 FREAS

NASA HQS

SS NEWELL, CLARK  
SG NAUGLE  
SCHARDT  
ROMAN  
SMITH  
DUBIN  
SL LIDDEL  
BRUNK  
FELLOWS  
HIPSHER  
HOROWITZ  
SM FOSTER  
ALLENBY  
GILL  
BADGLEY  
RR KURZWEG  
RTR NEILL  
ATSS SCHWIND [5]  
WX SWEET

OTHER CENTERSAMES R.C.

SONETT [5]  
LIBRARY [3]

LANGLEY

160 ADAMSON  
185 WEATHERWAX [2]  
213 KATZOFF  
DAVIS  
242 GARRICK  
235 SINGH

JPL

NEWBURN  
EIMER  
SNYDER  
SPEED  
CONEL  
KOVACH  
LOOMIS

UCLA

COLEMAN

UC BERKELEY

WILCOX